

11. JT, Davidow LS, Warshawsky D (1999) Tsix, a gene antisense to Xist at the X-inactivation centre. *Nat Genet* 17:400-404 [PubMed] [Cross Ref]10.1038/7734
12. Shen L, Wu LC, Sanfioglu S, Chen R, Mendoza AR, Dangel AW, Carroll MC, Zipf WB, Yu CY (1994) Structure and genetics of the partially duplicated gene RP located immediately upstream of the complement C4A and the C4B genes in the HLA class III region: molecular cloning, exon-intron structure, composite retroposon, and breakpoint of gene duplication. *J Biol Chem* 269:8466-8476 [PubMed]
13. Ostertag EM, Goodier JL, Zhang Y, Kazazian HH Jr (2003) SVA elements are nonautonomous retrotransposons that cause disease in humans. *Am J Hum Genet* 73:1444-1451 [PMC free article] [PubMed]
14. Wilund KR, Yi M, Campagna F, Arca M, Zuliani G, Fellin R, Ho YK, Garcia JV, Hobbs HH, Cohen JC (2002) Molecular mechanisms of autosomal recessive hypercholesterolemia. *Hum Mol Genet* 11:3019-3030 [PubMed] [Cross Ref]10.1093/hmg/11.24.3019
15. Kobayashi K, Nakahori Y, Miyake M, Matsumura K, Kondo-Iida E, Nomura Y, Segawa M, Yoshioka M, Saito K, Osawa M, et al (1998) An ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy. *Nature* 394:388-392 [PubMed] [Cross Ref]10.1038/28256
16. Gillics SD, Morrison SL, Oi VT, Tonegawa S (1983) A tissue-specific transcription enhancer element is located in the 5' region of the immunoglobulin heavy chain locus. *Cell* 33:717-728